arXiv:quant-ph/9706055v1 25 Jun 1997

Evaluation of vacuum energy for tensor fields on spherical spaces

E. N. Bukina[†] and N. N. Shtykov

Department of Theoretical Physics, The Irkutsk State University
[†]*N. N. Bogolubov LTF, JINR, 141980, Dubna, Russia*

31 May 1996

Abstract

The effective one-loop potential on $R^{m+1} \times S^N$ spaces for massless tensor fields is evaluated. The Casimir energy is given as a value of $\zeta-$ function by means of which regularization is made. In even- dimensional spaces the vacuum energy contains divergent terms coming from poles of $\zeta(s,q)$ at $s=1$, whereas in odd-dimensional spaces it becomes finite.

The appearance of divergent vacuum energy is one of peculiarities of quantum field theory. It is the well-known fact. However, in the presence of external constraints (external fields, nontrivial topology, curved space) the absolute value of vacuum energy, in principle, appears to be a measurble quantity. For instance, the phenomenon of nontrivial vacuum energy underlies the Casimir effect. Vacuum fluctuations in quantum field theory have been a subject of extensive research. [1] The contribution of vacuum energy in the Robertson-Walker model and Kaluza-Klein theory, where in place of an external field, the nontrivial background geometry (curved space) acts as the source to perturb vacuum, was computed in a series of works [2],[3], [4]. Analogous evaluations [5] are performed for massless scalar fields with coupling ξR and a massive spinor and vector fields on $R^m \times S^N$, $R^m \times CP^2$, where R is the curvature radius, R^m is the m-dimensional Euclidean space, S^N is the N-dimensional sphere, and CP^2 is the projective space.
In the present work, we have calculated the one-loop effective potential for massless tensor fields on $R^1 \times S^3$, $R^4 \times S^3$.

1. The One-Loop Potential for Tensor on $R^{m+1} \times S^N$

For the manifold $R^{m+1} \times S^N$ we introduce the metric as $g_{AB} = \eta_{AB} + r^2 h_{AB}$ where

$$\eta_{AB} = \left[\begin{array}{c|c} \eta_{ab} & 0 \\ \hline 0 & 0 \end{array}\right], \qquad h_{AB} = \left[\begin{array}{c|c} 0 & 0 \\ \hline 0 & h_{\alpha\beta} \end{array}\right]$$

$$A = 1, \ldots, m+1+N; \qquad a = 1, \ldots, m+1; \alpha = 1, \ldots, N$$

the equation for eigenvalues of the Lichnerowitz operator is as follows:

$$(\tilde{\Delta}T)_{AB} = -\nabla_c\nabla^c g_{AB}T^D_B - 2R_{ADBF}T^{DF} + R_{AD}T^D_B + R_{BD}T^D_A = \lambda T_{AB} \qquad (1)$$

On S^N, the curvature tensor and its convolutions are introduced by

$$R_{\alpha\beta\gamma\delta} = r^2[h_{\gamma\alpha}h_{\delta\beta} - h_{\alpha\delta}h_{\gamma\beta}], \quad R_{\beta\gamma} = (N-1)h_{\beta\gamma}, \quad R = N(N-1)/r^2. \qquad (2)$$

Let us expand the field considered harmonically as follows

$$T_{ab} = \bar{T_{ab}}(x)Y(y), \quad T_{a\alpha} = V_a(x)Y_\alpha(y), \quad T_{\alpha\beta} = S(x)Y_{\alpha\beta}. \qquad (3)$$

In terms of the metric tensor, expressions (2) and expansion (3), equation (1) takes the forms:

$$(\tilde{\Delta}T)_{ab} = (k^2+\lambda_0^2)T_{ab}, \quad (\tilde{\Delta}T)_{a\alpha} = (k^2+\lambda_1^2)T_{a\alpha}, \quad (\tilde{\Delta}T)_{\alpha\beta} = (k^2+\lambda_2^2)T_{\alpha\beta},$$

where

$$-\nabla_\alpha\nabla^\alpha Y = \lambda_0^2 Y, \quad (-\nabla_\beta\nabla^\beta + N - 1)Y_\alpha = \lambda_1^2 Y, \quad (-\nabla_\alpha\nabla^\alpha + 2N)Y_{\alpha\beta} = \lambda_2^2 Y_{\alpha\beta}.$$

Throughout we set r of sphere $= 1$, for simplicity.

1.1 Evaluation of eigenvalues λ_2^2 for tensor fields

The eigenvalues and degeneracies for scalar and tensor fields were determined in Ref. [5]. We consider the symmetric traceless second order tensor in the $(N+1)-$ dimensional Euclidean space $\hat{T_{\hat{\alpha}\hat{\beta}}}$ with

$$-\nabla_\alpha\nabla^\alpha\hat{T_{\hat{\alpha}\hat{\beta}}} = 0.$$

The symmetric fields on the sphere $S^N = SO(N+1)/SO(N)$ can be expressed through sperical harmonics on $SO(N+1)$ with the components in SO(N):

$$-\nabla^2 T_{\alpha\beta} = [c_2^{SO(N+1)}(l,p,0,0,\ldots) - c_2^{SO(N)}(2,0,0,\ldots)]T_{\alpha\beta}.$$

From the general theory of representations it follows that

$$c_2^{SO(N+1)}(l,p,0,0,\ldots) = l(l+N-2) + p(p+N-4),$$

which leads to

$$-\nabla^2 T_{\alpha\beta} = [l(l+N-1) + p(p+N-3) - 2N]T_{\alpha\beta}.$$

From the orthogonal expansion of tensor $T_{\alpha\beta}$ we obtain

$$T_{\alpha\beta} = T^\perp_{\alpha\beta} + \nabla_\alpha V^\perp_\beta + \nabla_\beta V^\perp_\alpha + (2 - \frac{2}{N})\nabla^2\phi$$

where $V_\alpha = V_\alpha^\perp + \nabla_\alpha \phi, \qquad \nabla^\alpha T^\perp_{\alpha\beta} = 0, \qquad \nabla^\alpha T^\perp_\alpha = 0.$ Therefore

$$\lambda^2_{2_s} = l(l+N-1), \quad \lambda^2_{2_\perp} = l(l+N-1)+2(N-1), \quad \lambda^2_{2_v} = l(l+N-1)+(N-2). \tag{4}$$

The degenerations of eigenvalues $\lambda^2_{2_\perp}$ are defined by dimensional representation $(l, 2, 0, \ldots)$

$$D_2(N,l) = \frac{(N+1)(N-2)(l+N)(l-1)(2l+N-1)(l+N-3)!}{2(N-1)!(l+1)!}. \tag{5}$$

The degenerations of $(\lambda^2_{2_\perp})_v$ and $(\lambda^2_2)_s$ are determined by dimensional representations $(l, 1, 0, 0 \ldots)$ and $(l, 0, 0, \ldots)$, respectively.

2.Evalution of the One-Loop Potential (OLP)

The initial object to calculate the effective potential is $T_{\mu\nu}(\tilde{\Delta}T)^{\mu\nu}$, where $\tilde{\Delta}$ was defined in Ref. [6]. The formal expression of OLP is given as

$$V^{(1)}_{eff} = \frac{1}{2\Omega(R^{m+1})} \text{Ln det}\tilde{\Delta},$$

where Ln det$\tilde{\Delta}$ is given by the following expression

$$\begin{aligned}
\text{Ln det}\tilde{\Delta} = \int d^{m+1}k\{\textstyle\sum_{n=0}^{\infty} \frac{(m+1)(m+2)}{2} D_0(N,n)\text{Ln}(\lambda_0^2 + k^2) + \\
+ \textstyle\sum_{n=1}^{\infty}(m+1)D_1(N,n)Ln(\lambda_1^2 + k^2) + \\
+ \textstyle\sum_{n=1}^{\infty}(m+1)D_0(N,n)Ln(\lambda_0^2 + k^2) + \sum_{n=2}^{\infty} D_1(N,n)Ln(\lambda_1^2 + k^2) + \\
+ \textstyle\sum_{n=2}^{\infty} D_2(N,n)Ln(\lambda_2^2 + k^2) + \sum_{n=2}^{\infty} D_0(N,n)Ln(\lambda_0^2 + k^2)\}
\end{aligned}$$

With $D_i(N,n)$ and λ_i^2 the regularized effective OLP has the form

$$\begin{aligned}
V^{(1)}_{tn} = Lim_{s\to 1} \tfrac{\Gamma(-s-m/2)\mu^{2\epsilon}}{2(4\pi)^{m/2}\Gamma(-s/2)} [\textstyle\sum_{n=0}^{\infty} \frac{(m+1)(m+2)}{2} \frac{(2n+N-1)\Gamma(n+N-1)}{\Gamma(N)n!} (n(n+N-1))^{\frac{m+s}{2}} + \\
+ \textstyle\sum_{n=1}^{\infty}(m+1) \frac{n(n+N-1)(2n+N-1)(n+N-3)!}{(N-2)!(n+1)!} (n(n+N-1) + (N-2))^{\frac{m+s}{2}} + \\
+ \textstyle\sum_{n=1}^{\infty}(m+1) \frac{(2n+N-1)\Gamma(n+N-1)}{\Gamma(N)n!} (n(n+N-1))^{\frac{m+s}{2}} + \\
+ \textstyle\sum_{n=2}^{\infty} \frac{n(n+N-1)(2n+N-1)(n+N-3)!}{(N-2)!(n+1)!} (n(n+N-1) + (N-2))^{\frac{m+s}{2}} + \\
+ \textstyle\sum_{n=2}^{\infty} \frac{(2n+N-1)(n+N-3)!(n+N)(N+1)(N-2)(n-1)}{2(N-1)!(n+1)!} (n(n+N-1) + 2(N-1))^{\frac{m+s}{2}} + \\
+ \textstyle\sum_{n=2}^{\infty} \frac{(2n+N-1)\Gamma(n+N-1)}{\Gamma(N)n!} (n(n+N-1))^{\frac{m+s}{2}}]
\end{aligned}$$

For $R^1 \times S^3$ representing the Robertson-Walker space-time, $V^{(1)}_{tn}(m=0, N=3) \approx -3.6563\frac{1}{\epsilon} - 4.2372$, where the relation

$$\sum_{n=0}^{\infty}(n+q)^l((n+q)^2 + M)^{\frac{m+s}{2}} = \sum_{n=0}^{\infty} \frac{\Gamma(k - \frac{m+s}{2})(-1)^k}{\Gamma(\frac{-m-s}{2})k!} \zeta(2k-l-m-s, q)M^k$$

is used. Also, we obtained for $R^4 \times S^3$: $V_{tn}^{(1)}(m=3, N=3) \approx 0.0262$. In this case the $\Gamma-$ function poles for negative values of the argument of $\Gamma(k-2)$ at $k=0,1,2$ are compensated by zero of $\zeta-$function at negative integer values of its argument according to the reflection property:

$$\Gamma(-z)\zeta(-2z) = \frac{\zeta(1+2z)\Gamma(\frac{1+2z}{2})}{\pi^{2z+1/2}}$$

with $\Gamma(-l) = \Gamma(-k)(-1)^{k+l}k!/l!$. It should be emphasized that the OLP on odd-dimensional manifolds has finite values. This is evident from the spectral properties of the Lichnerowicz operator that contrubutes only to the kinetic part of the Lagrangian, and the corresponding invariant counterterms R^2, $R^2_{\mu\nu}$, $R^2_{\mu\nu\sigma\delta}$ contain an even degree of r,only. Therefore they can't be constructed in an odd-dimensional space without some additional dimensional parameters. And vice versa, in an even-dimensional space the divergent part may be compensated by a specific combination of invariants constructed from the curvature tensor and contractions of the latter. In a $4-$ space this combination looks as $\alpha_1 R^2 + \alpha_2 R^2_{\alpha\beta} + \alpha_3 R^2_{\alpha\beta\gamma\delta}$ with arbitrary coefficients α_i. For the case of scalar and spinor fields this combination is defined as a conformal anomaly and turns out to be [6]

$$A_{(sc)} = \tfrac{1}{180} R_{\alpha\beta\gamma\sigma} R^{\alpha\beta\gamma\sigma} - \tfrac{1}{180} R_{\alpha\beta} R^{\alpha\beta} + \tfrac{1}{72} R^2,$$

$$A_{(sp)} = \tfrac{1}{720}\left[\tfrac{5}{2} R^2 - \tfrac{7}{2} R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma} - 4 R^{\mu\nu} R_{\mu\nu}\right].$$

The coefficients α_i were computed in a 4-dimensional space for the electromagnetic and gravitational fields [7]. The results demonstrate that the values of OLP for tensor fields give the main contribution to the vacuum energy in comparison with scalar, spinor and vector fields (compare with results of Ref. [5]). This means that the contributions to the vacuum energy on a compact space are ensured by the quantum corrections due to the gravitational field.

References

[1] *G. Plunien, B. Muller and W. Greiner,* Phys. Rep., **134** (1986) 87.

[2] *P. Candelas and S. Weinberg,* Nucl.Phys., **237** (1984) 397.

[3] *D. Birmingham and S. Sen,* Ann. Phys., **172** (1986) 451.

[4] *P. Chang and I. S. Dowker,* Nucl. Phys., **395** (1993) 407.

[5] *N. N. Shtykov,* Int. J. of Mod. Phys.A, vol. **10** (1995) 2333.

[6] *N. D. Birrell and P. C. W. Davies,* Quantum Field in Curved Space (Cambridge University Press,Cambridge,1982).

[7] *D. J. Toms,* Phys. Rev. **D27** (1983) 1803.